

# The Yang-Mills Field Strength Revisited

Samuel L. Marateck

*Courant Institute of Mathematical Sciences, New York University, New York, N.Y. 10012*

The Yang-Mills field strength incorporating a non-abelian feature is one of the cornerstones of the standard model. Although Yang-Mills gauge theories have been around for over fifty years, surprisingly the derivation of the Yang-Mills field strength using classical gauge theory does not appear anywhere in the literature. In their 1954 paper, Yang and Mills had to invent a non-Abelian field strength to satisfy certain criteria. In Section 5 we use Yang's gauge transformation in a heuristic derivation of the Yang-Mills field strength. The preceding sections cover material relating to the derivation. Section 3 shows where Pauli in the article cited by Yang and Mills gives an expression for the electro-magnetic field strength in terms of a commutator. For some reason, Yang and Mills did not use this approach.

## 1 INTRODUCTION

Although Yang-Mills (YM) gauge theory is now done using fiber-bundle theory – see, for instance, the review articles of Daniel [1] and Marateck [2] – it is of interest to analyse the way it developed using classical gauge theory.

In their seminal paper [5], when Yang and Mills discuss the phase factor - gauge transformation relationship, they cite Pauli's review paper [3]. It is interesting that, although Pauli in that paper presents the electromagnetic field strength in terms of a commutator, for whatever reason Yang and Mills did not use the commutator to obtain the Yang-Mills (YM) field strength – they obtained it by generalizing the electro-magnetic field strength. We provide an analysis of the steps required to derive the YM field strength in this way. Also presented is a derivation of this field strength using the commutator approach and the derivation of the YM field transformation using a slightly different way than is traditionally done.

## 2 GAUGE THEORY

Weyl [4] introduced as a phase factor an exponential in which the phase  $\alpha$  is preceded by the imaginary unit  $i$ , e.g.,  $e^{+iq\alpha(\mathbf{x})}$ , in the wave function for the wave equations (for instance, the Dirac equation is  $(i\gamma^\mu\partial_\mu - m)\psi = 0$ ). It is here that Weyl correctly formulated gauge theory as a symmetry principle from which electromagnetism could be derived. It had been shown that for a quantum theory of charged particles interacting with the electromagnetic field,

invariance under a gauge transformation of the potentials required multiplication of the wave function by the now well-known phase factor. Yang on page 19 of his selected papers [6] cites Weyl's gauge theory results as reported by Pauli [3] as a source for Yang-Mills gauge theory; although Yang didn't find out until much later that these were Weyl's results. Moreover, Pauli's article did not explicitly mention Weyl's geometric interpretation. It was only much after Yang and Mills published their article that Yang realized the connection between their work and geometry. In fact, on page 74 of his selected papers [6], Yang says

What Mills and I were doing in 1954 was generalizing Maxwell's theory. We knew of no geometrical meaning of Maxwell's theory, and we were not looking in that direction.

For the wave equations to be gauge invariant, i.e., have the same form after the gauge transformation as before, the local phase transformation  $\psi(\mathbf{x}) \rightarrow \psi(x)e^{+i\alpha(\mathbf{x})}$  has to be accompanied by the local gauge transformation

$$\mathbf{A}_\mu \rightarrow \mathbf{A}_\mu - q^{-1}\partial_\mu\alpha(\mathbf{x}) \tag{1}$$

This dictates that the  $\partial_\mu$  in the wave equations be replaced by the covariant derivative  $\partial_\mu + iqA_\mu$  in order for the  $\partial_\mu\alpha(\mathbf{x})$  terms to cancel each other. This pair of phase factor- gauge transformation is not unique. Another pair that retains gauge symmetry and results in the same covariant derivative has the  $q$  included in the phase factor, i.e.,  $\psi(\mathbf{x}) \rightarrow \psi(x)e^{+iq\alpha(\mathbf{x})}$  paired with

$$\mathbf{A}_\mu \rightarrow \mathbf{A}_\mu - \partial_\mu\alpha(\mathbf{x}) \tag{2}$$

The fact that this pairing is not unique is not surprising since a change in the phase factor and gauge transformation have no physical significance.

### 3 YANG-MILLS FIELD STRENGTH

Pauli, in equation 22a of Part I of his 1941 review article [3] gives the electromagnetic field strength in terms of a commutator<sup>1</sup>. In present-day usage it is

$$[D_\mu, D_\nu] = i\epsilon F_{\mu\nu} \tag{3}$$

where  $D_\mu$  is the covariant derivative  $\partial_\mu + i\epsilon A_\mu$ . Mathematically, equation (3) corresponds to the curvature (the field

---

<sup>1</sup>It is presented in Pauli's equation (22a)  $D_i D_k - D_k D_i = -i\epsilon f_{ik}$  where  $D_k = (\partial/\partial x_k) - i\epsilon\phi_k$ , and  $\phi_k$  is the electromagnetic potential,  $\epsilon$  is the charge, and  $f_{ik} = (\partial\phi_k/\partial x_i) - (\partial\phi_i/\partial x_k)$  is the field strength.

strength) reflecting the effect of parallel transport of a vector around a closed path, i.e., its holonomic behavior. If the field strength is zero, the vector will return to its point of origin pointing in its original direction. In their 1954 paper [5] Yang and Mills do not mention this relation, although they do cite Pauli's 1941 article [3]. They use

$$\psi = S\psi' \quad (4)$$

where  $\psi$  is a wave function and  $S$  is a local isotopic spin rotation represented by an  $SU(2)$  matrix, to obtain the gauge transformation in equation 3 of their paper

$$B'_\mu = S^{-1}B_\mu S + iS^{-1}(\partial_\mu S)/\epsilon \quad (5)$$

where  $B_\mu$  represents a field incorporating isotopic spin matrices. They<sup>2</sup> then define the field strength as

$$F_{\mu\nu} = (\partial_\nu B_\mu - \partial_\mu B_\nu) + i\epsilon(B_\mu B_\nu - B_\nu B_\mu) \quad (6)$$

They didn't know at that time that this corresponds to Cartan's second structural equation which in differential geometry notation is  $\Omega = d\mathbf{A} + [\mathbf{A}, \mathbf{A}]$ , where  $A$  is a connection on a principal fiber bundle.

On page 19 of his selected papers [6], Yang states

Starting from  $[\psi = S\psi'$  and  $S(\partial_\mu - i\epsilon B'_\mu)\psi' = (\partial_\mu - i\epsilon B_\mu)\psi]$  it was easy to get [our equation (5)]. Then I tried to define the field strength  $F_{\mu\nu}$  by  $F_{\mu\nu} = \partial_\nu B_\mu - \partial_\mu B_\nu$  which was a "natural" generalization of electromagnetism.

Yang returned to this work when he collaborated with Mills when they shared an office at Brookhaven. They published their results in their 1954 paper [5]. There they introduce equation (6) (their equation 4) by saying

In analogy to the procedure of obtaining gauge invariant field strengths in the electromagnetic case, we define now

$$(4)F_{\mu\nu} = (\partial_\nu B_\mu - \partial_\mu B_\nu) + i\epsilon(B_\mu B_\nu - B_\nu B_\mu)$$

One easily shows from  $[B'_\mu = S^{-1}B_\mu S + iS^{-1}(\partial_\mu S)/\epsilon]$  that

$$(5)F'_{\mu\nu} = S^{-1}F_{\mu\nu}S$$

under an isotopic gauge transformation. Other simple functions of  $B$  than (4) do not lead to such a simple transformation property.

---

<sup>2</sup>Yang had earlier started studying this problem as a graduate student at the University of Chicago and derived equation (5). When he returned to this problem as a visitor at Brookhaven, he in collaboration with Mills obtained (as we will explain) the field strength. See page 17 in Yang's selected papers, Ref. 3.

In section 5, we show that by substituting our equation (5) into the electromagnetic field strength  $F'_{\mu\nu} = \partial_\nu B'_\mu - \partial_\mu B'_\nu$  dictates adding a non-electromagnetic term, i.e., the non-Abelian term, so that their equation (5), the similarity transformation, is satisfied.

Using the Yang-Mills covariant derivative  $(\partial_\mu - i\epsilon B_\mu)$  let's see how the Yang-Mills field strength is obtained from the commutator

$$[D_\mu, D_\nu] = (\partial_\mu - i\epsilon B_\mu)(\partial_\nu - i\epsilon B_\nu) - (\partial_\nu - i\epsilon B_\nu)(\partial_\mu - i\epsilon B_\mu) \quad (7)$$

operating on the wave function  $\psi$ . Note that  $-\partial_\mu(B_\nu\psi) = -(\partial_\mu B_\nu)\psi - B_\nu\partial_\mu\psi$  and  $\partial_\nu(B_\mu\psi) = (\partial_\nu B_\mu)\psi + B_\mu\partial_\nu\psi$ . So we get a needed  $-B_\nu\partial_\mu$  and a  $B_\mu\partial_\nu$  term to cancel  $B_\nu\partial_\mu$  and  $-B_\mu\partial_\nu$  respectively. Thus expanding (7) we get

$$\begin{aligned} & \partial_\mu\partial_\nu - i\epsilon\partial_\mu B_\nu - i\epsilon B_\mu\partial_\nu - i\epsilon B_\nu\partial_\mu - \epsilon^2 B_\mu B_\nu - \partial_\nu\partial_\mu \\ & + i\epsilon\partial_\nu B_\mu + i\epsilon B_\nu\partial_\mu + i\epsilon B_\mu\partial_\nu + \epsilon^2 B_\nu B_\mu \end{aligned} \quad (8)$$

which reduces to  $i\epsilon(\partial_\nu B_\mu - \partial_\mu B_\nu) - \epsilon^2[B_\mu, B_\nu]$  or  $[D_\mu, D_\nu] = i\epsilon F_{\mu\nu}$

## 4 THE FIELD TRANSFORMATION

We present a detailed derivation of the gauge transformation by using the transformation

$$\psi' = S\psi \quad (9)$$

instead of the traditional  $\psi = S\psi'$ , i.e., the one Yang and Mills used. In order to obtain the gauge transformation in equation 3 of the Yang and Mills paper

$$B'_\mu = S^{-1}B_\mu S + iS^{-1}(\partial_\mu S)/\epsilon \quad (10)$$

requires you to use<sup>3</sup>  $\partial_\mu S^{-1} = -S^{-1}(\partial_\mu S)S^{-1}$ . Thus, the approach indicated by equation (9) is marginally more straight-forward since it doesn't require differentiating the inverse of a matrix. You must, however, perform this differentiation when deriving the field strength.

The covariant derivative,  $D_\mu = \partial_\mu - i\epsilon B_\mu$ , transforms the same way as  $\psi$  does

$$D'\psi' = SD\psi \quad (11)$$

---

<sup>3</sup>The following can be obtained by differentiating  $S^{-1}S = I$

The left-hand side of equation (11) becomes

$$(\partial_\mu - i\epsilon B'_\mu)S\psi = (\partial_\mu S)\psi + S\partial_\mu\psi - i\epsilon B'_\mu S\psi \quad (12)$$

But (12) equals  $S\partial_\mu\psi - i\epsilon SB_\mu\psi$ . Cancelling  $S\partial_\mu\psi$  on both sides we get,

$$(\partial_\mu S)\psi - i\epsilon B'_\mu S\psi = -i\epsilon SB_\mu\psi \quad (13)$$

or

$$B'_\mu S = SB_\mu + (\partial_\mu S)/(i\epsilon) \quad (14)$$

thus

$$B'_\mu = SB_\mu S^{-1} - i(\partial_\mu S)S^{-1}/\epsilon \quad (15)$$

We will use  $S = e^{i\alpha(\mathbf{x})\cdot\sigma}$ . So for  $\alpha$  infinitesimal,  $S = 1 + i\alpha \cdot \sigma$  which produces

$$\begin{aligned} B'_\mu &= (1 + i\alpha \cdot \sigma)B_\mu(1 - i\alpha \cdot \sigma) \\ &\quad - i(1/\epsilon)\partial_\mu(1 + i\alpha \cdot \sigma)(1 - i\alpha \cdot \sigma) \end{aligned} \quad (16)$$

Remembering that  $(a \cdot \sigma)(b \cdot \sigma) = a \cdot b + i\sigma \cdot (a \times b)$ , setting  $B_\mu = \sigma \cdot b_\mu$ , and since  $\alpha$  is infinitesimal, dropping terms of order  $\alpha^2$ , we get

$$\begin{aligned} b'_\mu \cdot \sigma &= b_\mu \cdot \sigma \\ &\quad + i[(\alpha \cdot \sigma)(b_\mu \cdot \sigma), (b_\mu \cdot \sigma)(\alpha \cdot \sigma)] + (1/\epsilon)\partial_\mu(\alpha \cdot \sigma) \end{aligned} \quad (17)$$

and finally

$$b'_\mu = b_\mu + 2(b_\mu \times \alpha) + (1/\epsilon)\partial_\mu\alpha \quad (18)$$

which (because our  $S$  is the inverse of Yang-Mills'  $S$ ) is equation 10 in the Yang-Mills paper [5].

## 5 FINDING THE FIELD STRENGTH

We reconstruct how one can go about determining the field strength. Since

$$F'_{\mu\nu} = S^{-1}F_{\mu\nu}S \quad (19)$$

let's start off with the electromagnetic-like field strength in the primed system

$$F'_{\mu\nu} = \partial_\nu B'_\mu - \partial_\mu B'_\nu \quad (20)$$

and express it in terms of the non-primed system fields. We calculate  $\partial_\nu B'_\mu$  from  $B'_\mu = S^{-1}B_\mu S + iS^{-1}(\partial_\mu S)/\epsilon$ , equation (5), obtaining

$$\begin{aligned} \partial_\nu B'_\mu &= -S^{-1}(\partial_\nu S)S^{-1}B_\mu S + S^{-1}(\partial_\nu B_\mu)S + S^{-1}B_\mu \partial_\nu S + \\ & i/\epsilon[-S^{-1}(\partial_\nu S)S^{-1}\partial_\mu S + S^{-1}\partial_\nu \partial_\mu S] \end{aligned} \quad (21)$$

So

$$\begin{aligned} \partial_\nu B'_\mu - \partial_\mu B'_\nu &= -S^{-1}[(\partial_\nu S)S^{-1}B_\mu - (\partial_\mu S)S^{-1}B_\nu]S \\ & + S^{-1}[\partial_\nu B_\mu - \partial_\mu B_\nu]S + S^{-1}[B_\mu \partial_\nu - B_\nu \partial_\mu]S + \\ & i/\epsilon[-S^{-1}(\partial_\nu S)S^{-1}\partial_\mu S + S^{-1}(\partial_\mu S)S^{-1}\partial_\nu S] \end{aligned} \quad (22)$$

We see that the  $+S^{-1}[\partial_\nu B_\mu - \partial_\mu B_\nu]S$  term satisfies equation (19) if the field strength only had the electromagnetic-like contribution. The other terms must either represent the transformed non-electromagnetic-like part of  $F_{\mu\nu}$  or be cancelled by adding the non-electromagnetic terms to equation (20). Since  $S$  is only used for the transformation, it should not appear in the expression for  $F_{\mu\nu}$ .

The  $i/\epsilon$  term in equations (22) dictates that a term multiplied by  $i\epsilon$  be added to equation (20). Since  $S^{-1}(\partial_\mu S)$  and  $S^{-1}\partial_\nu S$  appear in the expressions for  $B'_\mu$  and  $B'_\nu$  respectively, the product of  $S^{-1}(\partial_\mu S)$  and  $S^{-1}\partial_\nu S$  that appears in the last term of equation (22) suggests that we should start our quest to eliminate extra terms in equation (20) by adding  $i\epsilon B'_\mu B'_\nu$  to that equation. This product gives

$$\begin{aligned} i\epsilon B'_\mu B'_\nu &= i\epsilon[S^{-1}B_\mu S + iS^{-1}(\partial_\mu S)/\epsilon]* \\ & [S^{-1}B_\nu S + iS^{-1}(\partial_\nu S)/\epsilon] \end{aligned} \quad (23)$$

which equals

$$\begin{aligned}
& i\epsilon S^{-1} B_\mu B_\nu S - i/\epsilon S^{-1} (\partial_\mu S) S^{-1} \partial_\nu S - \\
& S^{-1} B_\mu \partial_\nu S - S^{-1} (\partial_\mu S) S^{-1} B_\nu S
\end{aligned} \tag{24}$$

All but the first term (which represents the transformation of  $i\epsilon B_\mu B_\nu$ ) cancel components of the extraneous terms in equation (22). And  $i\epsilon(B'_\mu B'_\nu - B'_\nu B'_\mu)$  cancels all of the extraneous terms except the transformation of  $i\epsilon(B_\mu B_\nu - B_\nu B_\mu)$ .

After performing the cancellation, we get

$$\begin{aligned}
& \partial_\nu B'_\mu - \partial_\mu B'_\nu + i\epsilon(B'_\mu B'_\nu - B'_\nu B'_\mu) = \\
& S^{-1} [\partial_\nu B_\mu - \partial_\mu B_\nu + i\epsilon(B_\mu B_\nu - B_\nu B_\mu)] S
\end{aligned} \tag{25}$$

which satisfies equation (19).

Using equation (15) instead of equation (10), the equation Yang and Mills used, does not simplify the generation of the Yang-Mills field strength.

As we know today, the commutator part of the YM field strength represents the interaction of the quanta of the  $B$  field, which is due to the isospin they carry.

## 6 CONCLUDING REMARK.

The conserved current of electromagnetism is associated with the U(1) gauge group which has the photon as its gauge particle. Because Yang and Mills worked with an SU(2) symmetry, their theory predicted three gauge particles. However, since the gauge invariance is local, there can be no mass term in the YM Lagrangian, so the gauge particles predicted by the YM theory have zero mass. This was corrected by the electroweak SU(2) x U(1) theory which incorporates YM non-abelian gauge theory. It predicts the existence of four gauge bosons: the three massive ones,  $W^\pm$  and  $Z^0$ , and the photon.

## ACKNOWLEDGEMENTS

The author thanks J. D. Jackson for his useful comments.

## REFERENCES

- [1] Daniel, M. and Viallet, C.M., *Rev. Mod. Physics.* **52** 175 (1980).
- [2] Marateck, Samuel L., *Notic. Amer. Math. Soc.* **53** 744 (2006); *Math. Advance in Translation, Chinese Acad. of Sciences (Chinese Translation)*, **2** 97, (2009).
- [3] Pauli, W., *Rev. Mod. Physics.* **13** 203 (1941).
- [4] Weyl, Hermann, *Zeit. f. Physic.* **330** 56 (1929).
- [5] Yang, C. N. and Mills, R. L., *Phys. Rev.* **96** 191 (1954).
- [6] Yang, C.N., *Selected Papers (1945-1980) With Commentary*, World Scientific (2005).